

University of Groningen

To the Editor-10-second ECG-based RMSSD as valid measure of HRV

Teegne, Balewgizie Sileshi; Man, Tengfei; van Roon, Arie M.; Riese, Harriette; Snieder, Harold

Published in:
Heart Rhythm

DOI:
[10.1016/j.hrthm.2018.10.038](https://doi.org/10.1016/j.hrthm.2018.10.038)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Final author's version (accepted by publisher, after peer review)

Publication date:
2019

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Teegne, B. S., Man, T., van Roon, A. M., Riese, H., & Snieder, H. (2019). To the Editor-10-second ECG-based RMSSD as valid measure of HRV. *Heart Rhythm*, 16(3), E35-E35.
<https://doi.org/10.1016/j.hrthm.2018.10.038>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy


If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

Accepted Manuscript

A 10-second ECG-based Root Mean Square of Successive Differences is a valid and reliable measure of heart rate variability

Balewgizie Sileshi Tegegne, MSc, Tengfei Man, MSc, Arie M. van Roon, PhD, Harriëtte Riese, PhD, Harold Snieder, PhD



PII: S1547-5271(18)31128-7

DOI: <https://doi.org/10.1016/j.hrthm.2018.10.038>

Reference: HRTM 7799

To appear in: *Heart Rhythm*

Received Date: 18 October 2018

Please cite this article as: Tegegne BS, Man T, van Roon AM, Riese H, Snieder H, A 10-second ECG-based Root Mean Square of Successive Differences is a valid and reliable measure of heart rate variability, *Heart Rhythm* (2018), doi: <https://doi.org/10.1016/j.hrthm.2018.10.038>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

A 10-second ECG-based Root Mean Square of Successive Differences is a valid and reliable measure of heart rate variability

Balewgizie Sileshi Tegegne, MSc^a; Tengfei Man, MSc^a; Arie M. van Roon, PhD^b; Harriëtte Riese, PhD^c; Harold Snieder, PhD^{a,*}

Departments of ^aEpidemiology, ^bVascular Medicine, ^cPsychiatry, University of Groningen, University Medical Center Groningen, The Netherlands.

*Correspondence: h.snieder@umcg.nl

We thank Lombardi et al.¹ for highlighting the relevance of our paper² in their editorial comment, but strongly disagree with their conclusion that the results of our study based on short-term heart rate variability (HRV) are “difficult to interpret”. We initially shared the same skepticism and concerns regarding the utility of ultra-short HRV measurements based on 10-second ECGs, which is why we set out to extensively test this in a validation study³ prior to using 10-second Root Mean Square of Successive Differences (RMSSD) in our recent publication². To validate, we assessed RMSSD from three separate 10-second recordings and gold standard measurements of 4 to 5 minutes and found substantial agreement ($r=0.853$ to 0.862) with only minimal deviation from the true mean, and short-term reproducibility for the 10-second recordings ranging from $r = 0.740$ to 0.751 .³ Thus, we conclude that a single 10-second ECG recording yields a valid RMSSD measurement with good reliability.

As such we do not believe, as stated by Lombardi et al.¹, that the negative findings for psychosocial factors were mainly caused by the use of ultra-short HRV measurements.

Furthermore, in contrast to Lombardi et al.’s view, we did consider instantaneous heart rate by additionally conducting all analyses on a measure of RMSSD adjusted for heart rate by the coefficient of variation, a parsimonious correction formula². Subsequently, our very large cohort study confirmed the well-known strong cross-sectional associations between RMSSD and antidepressant medication use, type-2 diabetes, and hypertension. Increased future application of RMSSD based on 10-second ECGs in prospective epidemiological studies may yield important new findings related to the development of cardiovascular morbidity and mortality.

References

1. Lombardi F, Huikuri H, Schmidt G, Malik M: Short-term heart rate variability: Easy to measure, difficult to interpret. *Heart Rhythm* 2018;15:1559–1560.
2. Tegegne BS, Man T, van Roon AM, Riese H, Snieder H: Determinants of heart rate

variability in the general population: The Lifelines Cohort Study. *Hear Rhythm* 2018;15:1552–1558.

3. Munoz ML, Van Roon A, Riese H, Thio C, Oostenbroek E, Westrik I, De Geus EJC, Gansevoort R, Lefrandt J, Nolte IM, Snieder H: Validity of (ultra-)short recordings for heart rate variability measurements. *PLoS One* 2015;10:e0138921.